

CONTRIBUTIONS
FROM THE
CUSHMAN LABORATORY
FOR
FORAMINIFERAL RESEARCH

INDEX TO VOLUMES 11-15

PRICE \$1.00 POSTPAID

SHARON, MASSACHUSETTS, U. S. A.

CONTRIBUTIONS FROM THE CUSHMAN
LABORATORY FOR FORAMINIFERAL RESEARCH

INDEX TO VOLUMES 11—15 INCLUSIVE

Owing to the demand of those who wish to bind these Contributions, the following Index has been prepared.

All new genera and species appear in bold-faced type, others in italic type. An abbreviated form gives the geologic formation and general locality with each species.

Papers are grouped both by geographic and geologic headings as well as by subjects.

In order to conserve space, the following abbreviated form has been used. Volume number with the part number following in parentheses followed by a colon. After the colon comes the page number, then the plate number in bold-faced type, and the figure number in italics. It is hoped that this will give ready reference to the subject matter of these volumes.

[Alabama]. Three New Rotaliform Foraminifera from the Lower Oligocene and Upper Eocene of Alabama, 14(3): 62. New Species of Foraminifera from the Lower Oligocene of Alabama, 15(3): 45. Eocene Foraminifera of Wilcox Age from Woods Bluff, Alabama, 15(4): 77.

Allomorphina, 12(4): 71; *allomorphinoides* (Cret.-Mexico), 12(4): 72; *contraria* (Cret.-Germany), 12(4): 72; *cretacea* (Cret.-Germany), 12(4): 71; **minuta** (Cret.-Texas), 12(4): 72, 13, 3; *navarroana* (Cret.-Texas), 12(4): 73, 13, 1; *obliqua* (Cret.-Germany), 12(4): 71; *trigona* (Cret.-Europe), 12(4): 72; *trochoides* (Cret.-Europe, America), 12(4): 71; *velascoensis* (Cret.-Mexico, Trinidad), 12(4): 72, 13, 2.

Allomorphinella contraria (Cret.-Europe), 12(4) : 73.

[America]. Notes on Some American Cretaceous Flabellinas,

11(4) : 83. Some American Cretaceous Buliminas, 11(4) : 96.

Notes on Some American Cretaceous Frondicularias, 12(1) : 11.

Some American Eocene Buliminas, 12(2) : 39. Some American

Cretaceous Species of Ellipsonodosaria and Chrysalogonium,

12(3) : 51. The Described American Eocene Species of Uvigerina,

13(3) : 74. A Few New Species of American Cretaceous

Foraminifera, 13(4) : 100. Additional New Species of American

Cretaceous Foraminifera, 14(2) : 31. Some New Species of

Rotaliform Foraminifera from the American Cretaceous,

14(3) : 66. New American Cretaceous Foraminifera, 15(4) : 89.

Ammobaculites cubensis (Eoc.-Cuba), 13(4) : 106, 16, 4, 16-18;

penonensis (Eoc.-Cuba), 13(4) : 107, 16, 13-15.

Ammocibicides pontoni (Eoc.-Ala.), 15(4) : 89.

Amphimorphina in the Eocene of Cuba, The Foraminiferal Genus,

12(1) : 1.

Amphimorphina crassa (Eoc.-Cuba), 12(1) : 3, 1, 10; *ignota*

(Eoc.-Calif.), 15(2) : 27, 6, 10-13; *lirata* (Eoc.-Cuba), 12(1) : 2,

1, 6-8; *polita* (Eoc.-Cuba), 12(1) : 2, 1, 9; *tenuistriata* (Eoc.-

Cuba), 12(1) : 1, 1, 1-3; var. *seminuda* (Eoc.-Cuba), 12(1) : 2, 1,

4, 5.

Amphistegina pinarensis (Eoc.-Cuba), 13(4) : 109, 16, 10-12.

Angulogerina, Notes on the Oligocene Species of Uvigerina and,

14(4) : 74.

Angulogerina byramensis (Olig.-Amer.), 14(4) : 87, 15, 18, 19;

elongata (Eoc.-France), 13(2) : 59, 8, 13; *europaea* (Eoc.-

France), 13(2) : 61, 8, 17, 18; *eximia* (Mioc.-Jamaica), 12(1) : 3,

1, 11, 12; *germanica* (Olig.-Germany), 14(4) : 85, 15, 14-16;

halkyardi (Eoc.-France), 13(2) : 60, 8, 14; *hispidula* (Olig.-

Ala.), 15(3) : 47, 9, 6; *muralis* (Eoc.-France), 13(2) : 55, 8, 3-5;

naranjoensis (Eoc.-Cuba), 13(1) : 16, 1, 56-58; *oligocaenica*

(Olig.-Germany), 14(4) : 86, 15, 8-11; var. *globosa* (Olig.-Ger-

many), 14(4) : 86, 15, 12, 13; *parvula* (Eoc.-Texas), 13(3) : 86,

12, 19; *pulchella* (Eoc.-France), 13(2) : 61, 8, 19; *rugoplicata*

(L. Olig.-Miss.), 11(2) : 33, 5, 5; (Olig.-Miss.), 14(4) : 88, 15,

20; *sagriniformis* (Olig.-Germany), 14(4) : 87, 15, 17; *tenui-*

striata (Olig.-Germany), 14(4) : 84, 15, 1-7; *vicksburgensis*

(L. Olig.-Miss.), 11(2) : 33, 5, 3, 4; (Olig.-Miss.), 14(4) : 88, 15,

21, 22; *wilcoxensis* (Eoc.-Ala.), 15(4) : 84, 14, 24, 25.

Anomalina affinis (Eoc.-Georges Bank), 15(3) : 75; *almendarensis*

(Eoc.-Cuba), 13(1) : 27, 2, 60, 61; *californiensis* (Tert.-Calif.),

- 11(3) : 64, 9, 8; *corrugata* (Eoc.-Cuba), 13(1) : 27, 2, 57-59; *crassisepta* (Eoc.-Calif.), 11(4) : 95, 14, 12; var. *caribaea* (Eoc.-Cuba), 12(3) : 62, 11, 12-14; *dorri*, var. *aragonensis* (Eoc.-Calif.), 15(2) : 32, 7, 5; *garzaensis* (Eoc.-Calif.), 15(2) : 32, 7, 3; *semicomplanata* (Cret.-Tenn.), 14(3) : 68, 12, 1.
- Articulina, and A New Genus Nodobaculariella, Notes on Some of the Species Referred to Vertebralina and, 13(2) : 41.
- Articulina byramensis*, 13(2) : 45; *compressa*, 13(2) : 45; *conico-articulata*, 13(2) : 45; *foveolata*, 13(2) : 45; *funalis*, 13(2) : 45; var. *inornata*, 13(2) : 45; *gibbula*, 13(2) : 45; *lineata*, 13(2) : 45; *mayori*, 13(2) : 45; *mexicana*, 13(2) : 45; *multilocularis*, 13(2) : 45; *nitida*, 13(2) : 45; *sagra*, 13(2) : 45; *sulcata*, 13(2) : 45; *terquemii*, 13(2) : 45.
- Asterigerina alabamensis* (Eoc.-Ala.), 15(4) : 86, 15, 11; *crassa-formis* (Eoc.-Calif.), 11(4) : 94, 14, 10; *wilcoxensis* (Eoc.-Ala.), 15(4) : 86, 15, 10.
- Astrononion, A New Genus of the Foraminifera, and Its Species, 13(1) : 29.
- Astrononion*, 13(1) : 30; *australe* (Olig.-Australia), 13(1) : 33, 3, 13, 14; *fijiense* (Rec.-Indo-Pac.), 13(1) : 35, 3, 15, 16; *italicum* (Plioc.-Italy), 13(1) : 35, 3, 19, 20; *novo-zealandicum* (Rec.-New Zeal.), 13(1) : 35, 3, 18; *sidebottomi* (Rec.-Medit.), 13(1) : 31, 3, 8; *stellatum* (Rec.-N. Atl.), 13(1) : 32, 3, 9-11; *stelligerum* (Rec.-Atl.), 13(1) : 31, 3, 7; *tumidum* (Rec.-S. Atl.), 13(1) : 33, 3, 17; *viragoense* (Rec.-E. Pac.), 13(1) : 32, 3, 12.
- Baggina marielina* (Eoc.-Cuba), 13(1) : 25, 2, 21-23.
- Bathysiphon* sp.(?) (Tert.-Calif.), 11(3) : 55, 8, 1.
- Bifarina atlantisae* (Eoc.-Georges Bank), 15(3) : 65, 11, 10; *nuttalli* (Eoc.-Calif.), 15(2) : 28, 6, 6; *tombigbeensis* (Eoc.-Ala.), 15(4) : 82, 14, 22, 23.
- Bifarinella*, 12(2) : 45; *ryukyuensis* (Pleist.-Japan), 12(2) : 46, 8, 7, 8.
- Bigenerina humblei* (Mioc.-La.), 15(1) : 3, 1, 4; *nodosaria*, var. *directa* (Mioc.-La.), 15(1) : 4, 1, 5.
- Bitubulogenerina howei*, A New Species from the Lower Oligocene, 11(1) : 20.
- Bitubulogenerina howei* (Olig.-Miss.), 11(1) : 20, 3, 10-12.
- Bolivinas from California, New Late Tertiary, 11(1) : 16.
- Bolivina atlantisae* (Eoc.-Georges Bank), 15(3) : 65, 11, 6, 7; *capdevilensis* (Eoc.-Cuba), 13(1) : 14, 1, 49, 50; *cochei* (Tert.-Calif.), 11(1) : 19, 3, 6, 7; *costata*, var. *dissimilis* (Mioc.-La.),

- 15(1): 6, 1, 10; cf. *decurrens* (Cret.-Calif.), 11(3): 73, 11, 9; *eggeri* (Cret.-Europe), 14(1): 28; *garretti* (L. Olig.-Miss.), 11(2): 33, 5, 2; *harangensis* (Mioc.-La.), 15(1): 6, 1, 9; *incrassata* (Cret.-Calif.), 11(3): 73, 11, 10; *marginata*, var. *gracillima* (Mioc.-Calif.), 14(1): 29; *marielina* (Eoc.-Cuba), 12(2): 29, 5, 10; *mornhinvegi* (L. Olig.-Miss.), 11(2): 32, 5, 1; *palmerae* (Eoc.-Cuba), 12(2): 29, 5, 13; *scabrata* (Eoc.-Cuba), 12(2): 29, 5, 11, 12; *sinuata*, var. *alisoensis* (Tert.-Calif.), 11(1): 19, 3, 5; *subadvena*, var. *sulphurensis* (Tert.-Calif.), 11(1): 20, 3, 8, 9; *submarina* (Eoc.-Georges Bank), 15(3): 64, 11, 8, 9; *tenuistriata* (Mioc.-La.), 15(1): 5, 1, 8; *tricosta* (Eoc.-Cuba), 13(1): 14, 1, 51, 52; sp. (?) (Tert.-Calif.), 11(3): 62, 9, 5.
- Bolivinella alata* (Eoc.-Cuba), 13(1): 12, 1, 43; *cubensis* (Eoc.-Cuba), 13(1): 13, 1, 46.
- Bolivinita costifera* (Cret.-Ark.), 13(4): 105, 15, 15.
- Bolivinoides austinana* (Cret.-Texas), 13(4): 104, 15, 10; *texana* (Cret.-Texas), 13(4): 104, 15, 12.
- Buliminas, Some American Cretaceous, 11(4): 96. Some American Eocene Buliminas, 12(2): 39. Notes on Some Oligocene Species of Bulimina and Buliminella, 13(1): 36. Notes on Some European Eocene [Miocene] Species of Bulimina, 13(2): 46. Notes on Some of the Early Described Eocene Species of Bulimina and Buliminella, 13(3): 65. Notes on Some Pliocene and Pleistocene Species of Bulimina and Buliminella, 14(3): 53. The Recent Species of Bulimina Named by d'Orbigny in 1826, 14(4): 90. Bulimina macilenta Cushman and Parker, A New Name, 15(4): 93.
- Bulimina acanthia*, 14(3): 61; *acicula*, 14(3): 61; *aculeata* (Rec.-Medit.), 14(4): 92, 16, 8-10; *acuta*, 14(3): 61; *adamsi* (Eoc.-Calif.), 12(2): 41, 7, 6; *affinis*, var. *striatula*, 14(3): 62; var. *tenuissimestriata* (Mioc.-Austria), 13(2): 53; *alsatica* (Olig.-France), 13(1): 39, 4, 6, 7; *andreaei* (Mioc.-Hungary), 13(2): 52; *arcuata* (Mioc.-France), 13(2): 53, 13(3): 72; *ariminensis* (Rec.-Medit.), 14(4): 92, 16, 11; *arkadelphiana* (Cret.-Amer.), 11(4): 96, 15, 1, 2; var. *midwayensis* (Eoc.-Texas), 12(2): 42, 7, 9, 10; *auriculata*, 13(3): 73; *brevis* (Rec.-Medit.), 14(4): 93, 16, 15; *buccinoides* (Mioc.-Germany), 13(2): 52; *buchiana* (Mioc.-Austria), 13(2): 48, 6, 6, 7; var. *calabra* (Mioc.-Hungary), 13(2): 51, 6, 8; *bulbiformis* (Mioc.-Italy), 13(2): 52; *cacumenata* (Eoc.-Texas), 12(2): 40, 7, 3; *calcarata* (Mioc.-Italy), 13(2): 52; *candida*, 13(3): 72; *caudigera* (Rec.-Medit.), 14(4): 94, 16, 17, 18; *conulus*, 13(3): 73; *convoluta*,

var. *dehiscens* (Mioc.-Australia), 13(2) : 53; *coprolithoides* (Olig.-France), 13(1) : 37, 4, 2; *corrugata* (Eoc.-Calif.), 11(4) : 92, 14, 7; (Eoc.-Georges Bank), 15(3) : 64; *costata* (Plioc.-Europe), 14(3) : 53, 9, 1, 2; *curtissima* (Eoc.-Calif.), 11(4) : 93, 14, 9; *cuspidata* (Mioc.-Hungary), 13(2) : 52; *cylindracea*, 14(3) : 61; *decorata*, 13(3) : 73; *denticulata* (Eoc.-Calif.), 12(2) : 42, 7, 7, 8; (Eoc.-Georges Bank), 15(3) : 64; *echinata* (Plioc.-Europe), 14(3) : 54, 9, 3, 4; *elegans* (Rec.-Medit.), 14(4) : 93, 16, 13; var. *gibba* (Mioc.-Austria), 13(2) : 53; *ellipsoides*, 14(3) : 61; *elongata* (Mioc.-Europe), 13(2) : 49, 7, 1-3; (Rec.-Medit.), 14(4) : 93, 16, 12; var. *cylindrica*, 13(3) : 73; var. *lappa* (Mioc.-Austria), 13(2) : 51, 7, 8; var. *subulata* (Mioc.-Austria), 13(2) : 51, 7, 6, 7; var. *tenera* (Mioc.-Europe, Egypt), 13(2) : 50, 7, 5; *eocena*, 13(3) : 72; *etnea* (Plioc.-Italy), 14(3) : 59, 10, 6-9; *excavata* (Eoc.-Calif.), 12(2) : 41, 7, 4; *exigua* (Cret.-Amer.), 11(4) : 99, 15, 7; *fossa* (Plioc.-Calif.), 14(3) : 56, 9, 10; *garzaensis* (Eoc.-Calif.), 11(4) : 93, 14, 8; *glanduliformis*, 13(3) : 72; *guayabalensis*, var. *ampla* (Eoc.-Calif.), 12(2) : 43, 8, 1; *imbricata*, var. *procera*, 13(3) : 73; *incrassata* (Mioc.-Hungary), 13(2) : 52; *inflata* (Plioc.-Europe), 14(3) : 58, 10, 4, 5; *instabilis* (Eoc.-Calif.), 12(2) : 44, 8, 3; *intorta*, 13(3) : 73; *jarvisi* (Eoc.-Trinidad), 12(2) : 39, 7, 1; *jugosa* (Olig.-Ecuador), 13(1) : 38, 4, 5; *laevigata* (Rec.-Medit.), 14(4) : 94, 16, 16; *lirata* (Eoc.-Calif.), 12(2) : 43, 8, 2; *longiscata*, 13(3) : 72; *macilenta* (Eoc.-Calif.), 15(4) : 93; *madagascarensis* (Rec.-Indian Ocean), 14(4) : 94, 16, 19, 20; *mammillata*, 14(3) : 61; *marginata* (Rec.-Medit.), 14(4) : 91, 16, 5, 6; *marginospinata* (Plioc.-Calif.), 14(3) : 57, 9, 11; *microcostata* (Eoc.-Calif.), 12(2) : 39, 7, 2; *minutissima*, 14(3) : 62; *obliqua*, 13(3) : 73; *obscura*, 13(3) : 72; *obtusa* (Cret.-Calif.), 11(3) : 73; *ornata* (Cret.-Amer.), 11(4) : 97, 15, 4; *ovata* (Mioc.-Austria), 13(2) : 47, 6, 4, 5; var. *apiculata* (Plioc.-Italy), 14(3) : 55, 9, 5; *oviformis*, 13(3) : 72; *ovigera*, 13(3) : 72; *ovula*, 13(3) : 73; *pagoda*, var. *deformata* (Plioc.-Calif.), 14(3) : 58, 10, 3; var. *denudata* (Plioc.-Calif.), 14(3) : 57, 10, 1, 2; var. *hebespinata* (Plioc.-Calif.), 14(3) : 55, 9, 6, 7; *parvula* (Mioc.-Hungary), 13(2) : 52; *pedunculata*, 14(3) : 62; "*peucetiae affinis*", 14(3) : 62; *pitecusana*, 14(3) : 62; *porrecta* (Mioc.-Hungary), 13(2) : 52; *prolixa* (Cret.-Amer.), 11(4) : 98, 15, 5; *punctata* (Rec.-Medit.), 14(4) : 93, 16, 14; *pupa*, 13(3) : 73; *pupoides* (Mioc.-Austria), 13(2) : 47, 6, 2, 3; *pupula* (Eoc.-New Zeal.), 13(3) : 66, 9, 1, 2; *pustulosa*, 14(3) : 62; *pygmaea* (Mioc.-

Germany), 13(2) : 52; *pyrula* (Mioc.-Austria), 13(2) : 46, 6, 1; *quadrata* (Cret.-Amer.), 11(4) : 100, 15, 12-16; *reussi* (Cret.-Amer.), 11(4) : 99, 15, 8, 10; var. *navarroensis* (Cret.-Amer.), 11(4) : 100, 15, 11; *rhomboidalis*, 14(3) : 62; cf. *rinconensis* (Tert.-Calif.), 11(3) : 62, 9, 4; *rotula* (Mioc.-Austria), 13(2) : 53; *rudita* (Cret.-Amer.), 12(2) : 45; *scalariformis*, 13(3) : 73; *sculptilis* (Olig.-Miss.), 13(1) : 37, 4, 3; var. *laciniata* (Olig.-Wash.), 13(1) : 38, 4, 4; (Olig.-Oregon), 13(2) : 53; *selseyensis*, 13(3) : 73; *semistriata*, 14(3) : 61; *simplex* (Eoc.-France), 13(3) : 67, 9, 4; *socialis* (Tert.-Calif.), 11(3) : 62, 9, 2; (Olig.-Germany), 13(1) : 36, 4, 1; *spinata* (Cret.-Calif.), 11(3) : 72, 11, 11; *spinosa*, 14(3) : 62; *splendens*, 13(3) : 73; *stalacta* (Eoc.-Calif.), 12(2) : 42, 7, 5; *striata* (Rec.-Atl.), 14(4) : 90, 16, 1-3; *subacuminata* (Plioc.-Calif.), 14(3) : 56, 9, 9; *subcalva* (Plioc.-Calif.), 14(3) : 55, 9, 8; *sulcata* (Rec.-Medit.), 14(4) : 91, 16, 4; *tarda* (Eoc.-Georges Bank), 15(3) : 64; *taylorensis* (Cret.-Amer.), 11(4) : 96, 15, 3; *tenuistriata* (Eoc.-France), 13(3) : 70, 10, 6; *textilariformis*, 13(3) : 72; *triangularis* (Cret.-Amer.), 11(4) : 97, 15, 6; *trigona* (Eoc.-France), 13(3) : 67, 9, 5; *trihedra* (Cret.-Amer.), 11(4) : 100, 15, 9; *trilobata* (Rec.-Medit.), 14(4) : 92, 16, 7; *triquetra* (Mioc.-Hungary), 13(2) : 52; *trocheata*, 13(3) : 72; *truncana* (Eoc.-Germany), 13(3) : 66, 9, 3; var. *angusta*, 13(3) : 73; *tuberculata* (Mioc.-Germany), 13(2) : 50, 7, 4; *uviformis*, 13(3) : 72.

Buliminella and *Neobulimina*, Notes on Some Cretaceous Species of, 12(1) : 5. Notes on Some Oligocene Species of *Bulimina* and *Buliminella*, 13(1) : 36. Notes on Some of the Early Described Eocene Species of *Bulimina* and *Buliminella*, 13(3) : 65. Notes on Some Pliocene and Pleistocene Species of *Bulimina* and *Buliminella*, 14(3) : 53.

Buliminella apiculata (Olig.-Australia), 13(1) : 39, 4, 10; var. *hebetata* (Olig.-Australia), 13(1) : 40, 4, 11, 12; *bassendorfsensis* (Olig.-Oregon), 13(1) : 40, 4, 13; *carseyi* (Cret.-Texas), 12(1) : 8, 2, 6; var. *plana* (Cret.-Texas), 12(1) : 8, 2, 7; *curta*, var. *basispinata* (Plioc.-Calif.), 14(3) : 59, 10, 10; *cushmani* (Cret.-Texas), 12(1) : 8; *fabilis* (Cret.-Texas), 12(1) : 7, 2, 5; *grata* (Eoc.-Calif.), 15(2) : 27, 6, 14; *imbricata* (Cret.-Germany), 12(1) : 6, 2, 2; *irregularis* (Eoc.-France), 13(3) : 68, 9, 6, 7; *laevis* (Cret.-Germany), 12(1) : 6, 2, 3; *multicamera* (Plioc.-Italy), 14(3) : 60, 10, 11, 12; *obtusa* (Cret.-France), 12(1) : 6, 2, 1; *obtusata* (Olig.-Miss.), 13(1) : 39, 4, 8, 9; *pulchra* (Eoc.-France), 13(3) : 69, 10, 1, 2; *semi-nuda* (Eoc.-France),

13(3) : 70, 10, 4, 5; *striato-punctata* (Eoc.-France), 13(3) : 69, 10, 3; *terquemiana* (Eoc.-France), 13(3) : 71, 10, 7; *turbinata* (Eoc.-France), 13(3) : 68, 9, 8, 9; *vitrea* (Cret.-Miss.), 12(1) : 7, 2, 4.

[California]. New Late Tertiary Bolivinas from California, 11(1) : 16. A Foraminiferal Faunule from the Type San Lorenzo Formation, Santa Cruz County, California, 11(3) : 53. Cretaceous Foraminifera from the Moreno Shale of California, 11(3) : 65. A New Genus of Foraminifera, *Discorbinella*, from Monterey Bay, California, 11(4) : 89. New Species of Foraminifera from the Kreyenhagen Shale of Fresno County, California, 11(4) : 90. A New Siphogenerinoides from California, 12(4) : 91. A New Species of *Pulvinulinella* from the California Miocene, 14(1) : 1. *Rupertia* (?) *adamsi*, A New Species from the Pliocene of California, 15(2) : 21. Some New and Interesting Foraminifera from the Kreyenhagen Shale of California, 15(2) : 23. *Bulimina macilenta* Cushman and Parker, A New Name, 15(4) : 93.

Cancris cubensis (Eoc.-Cuba), 13(1) : 25, 2, 48-50.

Cassidulina arenasensis (Eoc.-Cuba), 12(2) : 37, 6, 14; *constellata* (Eoc.-Cuba), 12(2) : 35, 6, 10; *crassipunctata* (Tert.-Calif.), 11(3) : 63, 9, 10; *crispula* (Eoc.-Cuba), 12(2) : 35, 6, 8, 9; *globosa* (Eoc.-Georges Bank), 15(3) : 73; *havanensis* (Eoc.-Cuba), 12(2) : 36, 6, 11; *labiata* (Eoc.-Cuba), 12(2) : 37, 6, 12; sp. (?) (Tert.-Calif.), 11(3) : 63.

Chilostomella ovoidea (Cret.-Austria), 12(4) : 73.

Chilostomelloides cyclostoma (Cret.-Austria), 12(4) : 73.

Chrysalidinella cubana (Eoc.-Cuba), 13(1) : 15, 1, 53-55.

Chrysalogonium, Some American Cretaceous Species of *Ellipsonodosaria* and, 12(3) : 51.

Chrysalogonium dickersoni (Eoc.-Cuba), 12(2) : 28, 5, 6, 7; *eximium* (Cret.-Texas), 14(2) : 42, 7, 5-8; *laeve* (Eoc.-Cuba), 12(2) : 27, 5, 1, 2; *tenuicostatum* (Eoc.-Cuba), 12(2) : 27, 5, 3-5; *texanum* (Cret.-Texas), 12(3) : 55, 9, 24, 25.

Cibicides berryi (Cret.-Tenn.), 14(3) : 71, 12, 6; *carstensi* (Mioc.-La.), 15(1) : 13, 2, 8; var. *opima* (Mioc.-La.), 15(1) : 14, 2, 7; *cookei* (L. Olig.-Ala.), 14(3) : 65, 11, 3; *cushmani* (Eoc.-Calif.), 15(2) : 33, 7, 7; (Eoc.-Georges Bank), 15(3) : 76, 12, 26; *havanensis* (Eoc.-Cuba), 13(1) : 28, 3, 1-3; *lobatulus* (Eoc.-Georges Bank), 15(3) : 76, 12, 25; *martinezensis* (Eoc.-Calif.), 15(2) : 33, 7, 6; *pippeni* (L. Olig.-Ala.), 14(3) : 64, 11, 2; *praecursorius*

- (Eoc.-Ala.), 15(4): 88; *pseudoungerianus* (Eoc.-Georges Bank), 15(3): 76, 12, 23, 24; (Eoc.-Ala.), 15(4): 88, 15, 25, 26; var. *evolutus* (Tert.-Calif.), 11(3): 64, 9, 11; *stephensoni* (Cret.-Ala.), 14(3): 70, 12, 5; *venezuelanus* (Eoc.-Calif.), 15(2): 32, 7, 4; sp. (Eoc.-Ala.), 15(4): 88, 15, 23, 24.
- Clavulinoides cubensis* (Eoc.-Cuba), 13(1): 2, 1, 5-7; *eucarinatus* (Eoc.-Cuba), 13(1): 3, 1, 10, 11; *excurrens* (Eoc.-Cuba), 13(1): 3, 1, 14, 15; *havanensis* (Eoc.-Cuba), 13(1): 3, 1, 12, 13; *marielinus* (Eoc.-Cuba), 13(1): 4, 1, 16, 17; *subulatus* (Eoc.-Cuba), 13(1): 2, 1, 8, 9.
- Coleites reticulosus* (Eoc.-Ala.), 15(4): 87, 15, 14-20.
- Cornuspira byramensis* (L. Olig.-Miss.), 11(2): 27, 4, 4; (Eoc.-Ala.), 15(4): 79, 13, 8.
- [Cretaceous]. New Cretaceous Textulariidae, 11(1): 1. Cretaceous Foraminifera from the Moreno Shale of California, 11(3): 65. Notes on Some American Cretaceous Flabellinas, 11(4): 83. Some American Cretaceous Buliminids, 11(4): 96. Notes on Some Cretaceous Species of Buliminella and Neobulimina, 12(1): 5. Notes on Some American Cretaceous Frondicularias, 12(1): 11. Some American Cretaceous Species of Ellipsonodosaria and Chrysalogonium, 12(3): 51. Cretaceous Foraminifera of the Family Chilostomellidae, 12(4): 71. Some Notes on Cretaceous Species of Marginulina, 13(4): 91. A Few New Species of American Cretaceous Foraminifera, 13(4): 100. Cretaceous Species of Gümbelina and Related Genera, 14(1): 2. Additional New Species of American Cretaceous Foraminifera, 14(2): 31. Some New Species of Rotaliform Foraminifera from the American Cretaceous, 14(3): 66. New American Cretaceous Foraminifera, 15(4): 89.
- [Cribroglorotalia]. Additional New Species of Foraminifera, and a New Genus from the Eocene of Cuba, 12(3): 55.
- Cribroglorotalia*, 12(3): 63; *marielina* (Eoc.-Cuba), 12(3): 63, 11, 17-19.
- Cribrolinoides*, A New Genus of the Foraminifera, Its Development and Relationships, 15(1): 15.
- Cribrolinoides*, 15(1): 15; *albatrossi* (Rec.-Pac.), 15(1): 18; *curta* (Plioc.-Asia), 15(1): 17.
- [Cuba]. The Foraminiferal Genus Amphimorphina in the Eocene of Cuba, 12(1): 1. New Genera and Species of Foraminifera from the Eocene of Cuba, 12(2): 27. Additional New Species of Foraminifera and a New Genus from the Eocene of Cuba, 12(3): 55. Further New Species of Foraminifera from the

- Eocene of Cuba, 13(1): 1. Additional New Species of Eocene Foraminifera from Cuba, 13(4): 106.
- Cyclammina* cf. *clarki* (Tert.-Calif.), 11(3): 55, 8, 3; cf. *incisa* (Tert.-Calif.), 11(3): 55, 8, 4.
- Cycloloculina cubensis* (Eoc.-Cuba), 12(3): 61, 11, 15, 16.
- Darbyella wilcoxensis* (Eoc.-Ala.), 15(4): 79, 13, 11, 12.
- Dentalina angusticostata* (Cret.-Texas), 14(2): 41, 6, 21, 22; *basiplanata* (Cret.-Texas, Ark.), 14(2): 38, 6, 6-8; *basitorta* (Cret.-Miss.), 14(2): 37, 6, 4, 5; *communis*(?) (Eoc.-Ala.), 15(4): 80, 14, 1-3; *delicatula* (Cret.-Texas), 14(2): 40, 6, 19, 20; *havanensis* (Eoc.-Cuba), 13(1): 11, 1, 39, 40; *hispidocostata* (Eoc.-Calif.), 15(2): 25, 6, 4, 5; *involvens* (Cret.-Texas), 14(2): 37, 6, 3; cf. *megapolitana* (Cret.-Calif.), 11(3): 71, 10, 3; *nasuta* (Eoc.-Georges Bank), 15(3): 57, 10, 10, 11; *pertinens* (Cret.-Texas, Miss.), 14(2): 40, 6, 15-18; *pseudoinvolvens* (Olig.-Ala.), 15(3): 45, 9, 1, 2; cf. *pungens* (Eoc.-Georges Bank), 15(3): 56, 10, 9; cf. *soluta* (Eoc.-Georges Bank), 15(3): 56, 10, 1; *solvata* (Cret.-Miss.), 14(2): 39, 6, 9-14; sp.(?) A (Eoc.-Georges Bank), 15(3): 57, 10, 2; sp.(?) B (Eoc.-Georges Bank), 15(3): 57, 10, 3, 4; sp.(?) C (Eoc.-Georges Bank), 15(3): 57, 10, 5, 6; sp.(?) D (Eoc.-Georges Bank), 15(3): 57, 10, 7, 8.
- Discorbinella*, from Monterey Bay, California, A New Genus of Foraminifera, 11(4): 89.
- Discorbinella*, 11(4): 89; *montereyensis* (Rec.-Calif.), 11(4): 89, 14, 13.
- Discorbis arcuato-costata* (L. Olig.-Miss.), 11(2): 38, 5, 9; *cocoaensis* (Eoc.-Ala.), 14(3): 63, 11, 1; *havanensis* (Eoc.-Cuba), 13(1): 19, 2, 15, 16; *incrustedatus* (Eoc.-Cuba), 13(1): 21, 2, 17, 18; *subaraucana*, var. *dissona* (Olig.-La.), 15(1): 10, 2, 2; *subglobosa* (L. Olig.-Miss.), 11(2): 38, 5, 8; *subpatelliformis* (Olig.-Ala.), 15(3): 48, 9, 8.
- Dorothia principensis* (Eoc.-Cuba), 12(3): 57, 10, 3, 4; (Eoc.-Calif.), 15(2): 24, 6, 23.
- Ehrenbergina bicornis*, var. *eocenica* (Eoc.-Cuba), 12(2): 37, 6, 6; *cristata* (Eoc.-Cuba), 12(2): 38, 6, 15; *cubensis* (Eoc.-Cuba), 12(2): 38, 6, 13; *spinosissima* (Mioc.-Jamaica), 12(1): 5, 1, 15, 16.
- Ellipsoglandulina cubensis* (Eoc.-Cuba), 13(1): 17, 1, 66-68; *principiensis* (Eoc.-Cuba), 13(1): 18, 2, 1-3.

Ellipsolagena sculpturata (Eoc.-Cuba), 12(3) : 61, 11, 10, 11.

Ellipsonodosaria and *Chrysalogonium*, Some American Cretaceous Species of, 12(3) : 51.

Ellipsonodosaria alexanderi (Cret.-Texas), 12(3) : 52, 9, 6-9; var. **impensia** (Cret.-Texas, Ark.), 14(2) : 48, 8, 4, 5; **annulifera** (Eoc.-Cuba), 12(2) : 28, 5, 8, 9; (Eoc.-Georges Bank), 15(3) : 69; **atlantisae** (Eoc.-Georges Bank), 15(3) : 70, 12, 3, 4; var. **hispidula** (Eoc.-Georges Bank), 15(3) : 70, 12, 5; **cocodaensis** (Eoc.-Georges Bank), 15(3) : 68, 11, 27-33; **curvatura** (Eoc.-Georges Bank), 15(3) : 71, 12, 6; var. **spinea** (Eoc.-Georges Bank), 15(3) : 71, 12, 7-11; **dentata-glabrata** (Cret.-Texas), 12(3) : 54, 9, 22, 23; **exilis** (Cret.-Texas), 12(3) : 51, 9, 1, 2; (?) **granti** (Cret.-Texas, Ark.), 12(3) : 51, 9, 3-5; **horridens** (Cret.-Tenn.), 12(3) : 53, 9, 19-21; (?) **jarvisi** (Cret.-Trinidad), 12(3) : 53, 9, 16-18; **minuta** (Cret.-Texas), 14(2) : 48, 8, 6; **modesta**, var. **prolata** (Eoc.-Cuba), 13(4) : 109, 16, 1, 7; **pseudoscripta** (Cret.-Texas), 13(4) : 103, 15, 14; **semijugosa** (Eoc.-Georges Bank), 15(3) : 69, 12, 1, 2; **stephensoni** (Cret.-Texas), 12(3) : 52, 9, 10-15; var. **speciosa** (Cret.-Ark.), 14(2) : 47, 8, 3.

Elphidiella, 12(4), 89.

Elphidium and Related Genera, Some New Species of, 12(4) : 78.

Elphidium chapmani (Mioc.-Australia), 12(4) : 80, 14, 6; **crassatum** (Olig.-Australia), 12(4) : 81, 14, 8; **crespinae** (Olig.-Australia), 12(4) : 78, 14, 1; **culebreense** (Olig.-Panama), 12(4) : 80, 14, 4; **earlandi** (Rec.-E. Atl.), 12(4) : 85, 15, 5; **georgianum** (Tert.-Georges Bank), 11(4) : 79, 12, 5; **hispidulum** (Rec.-Australia), 12(4) : 83, 14, 13; **howchini** (Olig.-Australia), 12(4) : 79, 14, 2; **indicum** (Rec.-India), 12(4) : 83, 14, 10; **lidoense** (Rec.-Medit.), 12(4) : 86, 15, 6; **nigareense** (Rec.-Medit.), 12(4) : 85, 15, 3; **novo-zealandicum** (Rec.-New Zeal.), 12(4) : 85, 15, 4; **panamense** (Olig.-Panama), 12(4) : 79, 14, 3; **papillosum** (Rec.-Australia), 12(4) : 83, 14, 12; **parri** (Mioc.-Australia), 12(4) : 81, 14, 7; **pseudoinflatum** (Olig.-Australia), 12(4) : 80, 14, 5; **pseudonodosum** (Plioc.-Australia), 12(4) : 82, 14, 9; **rolshauseni** (Mioc.-La.), 15(1) : 4, 1, 7; **sculpturatum** (Rec.-Australia), 12(4) : 84, 15, 2; **subinflatum** (Mioc.-Australia), 12(4) : 84, 15, 1; **subplanatum** (Olig.-Germany), 12(4) : 82, 14, 11; sp. (?) (Eoc.-Georges Bank), 15(3) : 62.

Entosolenia byramensis (L. Olig.-Miss.), 11(2) : 31, 4, 10; **crumenata** (L. Olig.-Miss.), 11(2) : 31, 4, 9; (Eoc.-Georges Bank), 15(3) : 66, 11, 4, 5; cf. **laevis** (Eoc.-Georges Bank), 15(3) : 66, 11, 2; **marginata** (Eoc.-Georges Bank), 15(3) : 66, 11, 1, 3.

[Eocene]. New Species of Foraminifera from the Kreyenhagen Shale of Fresno County, California, 11(4) : 90. The Foraminiferal Genus Amphimorphina in the Eocene of Cuba, 12(1) : 1. New Genera and Species of Foraminifera from the Eocene of Cuba, 12(2) : 27. Some American Eocene Buliminas, 12(2) : 39. Additional New Species of Foraminifera and A New Genus from the Eocene of Cuba, 12(3) : 55. Further New Species of Foraminifera from the Eocene of Cuba, 13(1) : 1. Notes on the Early Described Eocene Species of Uvigerina and Some New Species, 13(2) : 54. Notes on Some of the Early Described Eocene Species of Bulimina and Buliminella, 13(3) : 65. The Described American Eocene Species of Uvigerina, 13(3) : 74. Additional New Species of Eocene Foraminifera from Cuba, 13(4) : 106. Three New Rotaliform Foraminifera from the Lower Oligocene and Upper Eocene of Alabama, 14(3) : 62. Some New and Interesting Foraminifera from the Kreyenhagen Shale of California, 15(2) : 23. Eocene Foraminifera from Submarine Cores off the Eastern Coast of North America, 15(3) : 49. Eocene Foraminifera of Wilcox Age from Woods Bluff, Alabama, 15(4) : 77. Bulimina macilenta Cushman and Parker, A New Name, 15(4) : 93.

Eponidella, 11(1) : 13; *libertadensis* (Mioc.-Venezuela), 11(1) : 14, 3, 1-4.

Eponides lotus (Eoc.-Ala.), 15(4) : 85, 15, 3-6; *marielensis* (Eoc.-Cuba), 13(1) : 23, 2, 27-29; *pricipiensis* (Eoc.-Cuba), 13(1) : 23, 2, 30-32; *pygmaeus* (Eoc.-Georges Bank), 15(3) : 72; *rutteni* (Eoc.-Cuba), 13(1) : 24, 2, 33-35; *umbonatus* (Eoc.-Georges Bank), 15(3) : 72.

[Europe]. Notes on Some European Eocene [Miocene] Species of Bulimina, 13(2) : 46.

Flabellinas, Notes on Some American Cretaceous, 11(4) : 83.

Flabellina jarvisi (Cret.-Amer.), 11(4) : 85, 13, 7, 8; *pilulifera* (Cret.-Calif.), 11(3) : 67, 10, 6; *reticulata* (Cret.-Amer.), 11(4) : 87, 13, 19; *rugosa* (Cret.-Amer.), 11(4) : 83, 13, 1-6; *suturalis* (Cret.-Amer.), 11(4) : 86, 13, 9-18.

Flintia laticoncava (L. Olig.- Miss.), 11(2) : 26, 4, 3.

Fronicularias, Notes on Some American Cretaceous, 12(1) : 11.

Fronicularia archiaciana (Cret.-Calif.), 11(3) : 70, 11, 8; (Cret.-Europe and Amer.), 12(1) : 19, 4, 8-10; *arkadelphiana* (Cret.-Ark.), 12(1) : 12, 3, 2, 3; *austinana* (Cret.-Texas), 12(1) : 13, 3, 12, 13; *clarki* (Cret.-Amer.), 12(1) : 12, 3, 4-6; *cordata*

(Cret.-Europe and Amer.), 12(1): 16, 4, 1-3; *cuspidata* (Cret.-Amer.), 12(1): 19, 4, 12-14; var. *costifera* (Cret.-Texas), 12(1): 21, 4, 15; *dimidia* (Cret.-Texas), 12(1): 11, 3, 1; *frankei* (Cret.-Europe and Amer.), 12(1): 18, 4, 6, 7; *garretti* (L. Olig.-Miss.), 11(2): 28, 4, 8; *glabrans* (Cret.-Tenn.), 12(1): 16; *goldfussi* (Cret.-Europe and Amer.), 12(1): 15, 3, 21, 22; *intermittens* (Cret.-Europe and Amer.), 12(1): 21, 4, 16-18; *inversa* (Cret.-Europe and Amer.), 12(1): 16, 3, 23, 24; *jarvisi* (Cret.-Trinidad), 15(4): 91, 16, 6; *linearis* (Cret.-Europe and Amer.), 12(1): 21, 4, 19, 20; *linguiformis* (Cret.-Europe and Amer.), 12(1): 17, 3, 18-20; *mucronata* (Cret.-Europe and Amer.), 12(1): 15, 3, 16, 17; *seminiformis* (Cret.-Calif.) 11(3): 70, 11, 5; *striatula* (Cret.-Europe and Amer.), 12(1): 18, 4, 4, 5; *undulosa* (Cret.-Texas), 12(1): 13, 3, 7-11; *verneuilliana* (Cret.-Europe and Amer.), 12(1): 19, 4, 11; *watersi* (Cret.-Amer.), 12(1): 14, 3, 14, 15; sp.(?) (Cret.-Calif.), 11(3): 70, 11, 6.

Gaudryina cubana (Eoc.-Cuba), 12(3): 56, 10, 2, 10, 11; *navarroana*, var. *crassaformis* (Cret.-Calif.), 11(3): 66, 10, 1; *subtenuis* (Rec.-Pac.), 14(1): 28; sp. (Eoc.-Ala.), 15(4): 78, 13, 3; (Eoc.-Ala.), 15(4): 78.

Gaudryina (*Pseudogaudryina*) *rutteni* (Eoc.-Cuba), 12(3): 56, 10, 15, 16.

Gaudryinella cubana (Eoc.-Cuba), 12(3): 55, 10, 1, 5, 6.

[Georges Bank]. Some New Foraminifera from the Late Tertiary of Georges Bank, 11(4): 77. Eocene Foraminifera from Submarine Cores off the Eastern Coast of North America, 15(3): 49.

Glandulina abbreviata (Eoc.-Ala.), 15(4): 81, 14, 10; *basi-spinata* (Tert.-Georges Bank), 11(4): 78, 12, 6-8; *cubensis* (Eoc.-Cuba), 13(1): 12, 1, 42; *laevigata* (Tert.-Calif.), 11(3): 59.

Globigerina altispira (Mioc.-Jamaica), 12(1): 5, 1, 13, 14; *dissimilis* (Eoc.-Cuba), 13(1): 25, 3, 4-6.

Globobulimina cf. *pacifica* (Tert.-Calif.), 11(3): 62, 9, 3.

Globorotalia centralis (Eoc.-Cuba), 13(1): 26, 2, 62-65; *crassata* (Eoc.-Georges Bank), 15(3): 74, 12, 19; var. *densa* (Eoc.-Georges Bank), 15(3): 74, 12, 20; *fohsi* (Mioc.-La.), 15(1): 12, 2, 6; *kochi* (Eoc.-Georges Bank), 15(3): 75, 12, 22; *mayeri* (Mioc.-La.), 15(1): 11, 2, 4; *palmerae* (Eoc.-Cuba), 13(1): 26, 2, 51-53; *quadraria* (Mioc.-La.), 15(1): 11, 2, 5; *spinulosa* (Eoc.-Georges Bank), 15(3): 75, 12, 21; *wilcoxensis* (Eoc.-Ala.), 15(4): 88, 15, 21, 22.

- Globotruncana cretacea* (Cret.-Tenn.), 14(3) : 67, 11, 6; (Cret.-Tenn.), 15(4) : 92, 16, 8.
- Globulina alabamensis* (Olig.-Ala.), 15(3) : 46, 9, 3; *inaequalis* (Eoc.-Georges Bank), 15(3) : 60, 10, 36; *münsteri* (Eoc.-Georges Bank), 15(3) : 61, 10, 38.
- Goëssella cubensis* (Eoc.-Cuba), 12(3) : 58, 10, 17, 18; *trinitatensis* (Eoc.-Georges Bank), 15(3) : 50, 9, 16.
- Gonatosphaera alternicostata* (Eoc.-Cuba), 12(2) : 31, 5, 16, 17; *principensis* (Eoc.-Cuba), 12(3) : 59, 11, 7-9.
- Grammostomum tessera*, 14(1) : 6, 1, 25.
- Gümbelina and Related Genera, Cretaceous Species of, 14(1) : 2.
- Gümbelina*, 14(1) : 3; *carinata* (Cret.-Texas), 14(1) : 18, 3, 10; *costulata* (Cret.-Texas), 14(1) : 16, 3, 7-9; *cubensis* (Eoc.-Georges Bank), 15(3) : 63, 10, 54; *excolata* (Cret.-Amer.), 14(1) : 17, 3, 11; *glabrans* (Cret.-Texas), 14(1) : 15, 3, 1, 2; *globocarinata* (Cret.-Texas), 14(1) : 10, 2, 4, 5; *globulosa* (Cret.-Europe and Amer.), 14(1) : 6, 1, 28-33; *lata* (Cret.-Germany), 14(1) : 12, 2, 11; *moremani* (Cret.-Texas), 14(1) : 10, 2, 1-3; *paucistriata* (Cret.-Texas), 14(1) : 18, 3, 12, 13; *planata* (Cret.-Texas), 14(1) : 12, 2, 13, 14; *plummerae* (Cret.-Amer.), 14(1) : 15, 3, 3-5; *pseudotessera* (Cret.-Texas), 14(1) : 14, 2, 19-21; *pulchra* (Cret.-Sweden), 14(1) : 12, 2, 12; *punctulata* (Cret.-Texas), 14(1) : 13, 2, 15, 16; *reussi* (Cret.-Texas), 14(1) : 11, 2, 6-9; *semicostata* (Cret.-Texas), 14(1) : 16, 3, 6; *spinifera* (Cret.-Tenn.), 14(1) : 11, 2, 10; *striata* (Cret.-Europe, Amer.), 14(1) : 8, 1, 34-40; *ultimatumida* (Cret.-Mexico), 14(1) : 13, 2, 17, 18; *venezuelana* (Eoc.-Georges Bank), 15(3) : 62, 10, 50-53.
- Gümbelitria*, 14(1) : 19; *cretacea* (Cret.-Amer.), 14(1) : 19, 3, 14; sp. (?) (Eoc.-Georges Bank), 15(3) : 63.
- Guttulina georgiana* (Tert.-Georges Bank), 11(4) : 78, 12, 4; *irregularis* (Eoc.-Georges Bank), 15(3) : 60, 10, 35; *wilcoxensis* (Eoc.-Ala.), 15(4) : 80, 14, 8, 9.
- Gyroidina arkadelphiana* (Cret.-Ark.), 14(2) : 49, 8, 7; *elongata* (Eoc.-Cuba), 13(1) : 22, 2, 36-38; *plummerae* (Eoc.-Cuba), 13(1) : 22, 2, 42-44; *protea* (Eoc.-Cuba), 13(1) : 22, 2, 39-41; *scalata* (Olig.-La.), 15(1) : 10, 2, 1; *soldanii* (Tert.-Calif.), 11(3) : 62, 9, 7; cf. *soldanii* (Eoc.-Georges Bank), 15(3) : 72, 12, 12-14.
- Haddonina cubensis* (Eoc.-Cuba), 13(4) : 107, 16, 8, 9.
- Hantkenina alabamensis* (Eoc.-Georges Bank), 15(3) : 74, 12, 18; cf. *dumblei* (Eoc.-Calif.), 15(2) : 32, 7, 2.

Heronallenia vicksburgensis (L. Olig.-Miss.), 11(2) : 38, 5, 11.

[Jamaica]. Three New Foraminifera from the Miocene, Bowden Marl, of Jamaica, 12(1) : 3.

Karrerriella arenasensis (Eoc.-Cuba), 13(1) : 5, 1, 20, 21; (Eoc.-Georges Bank), 15(3) : 51, 9, 14, 15; *cubensis* (Eoc.-Cuba), 13(1) : 4, 1, 18, 19; *halkyardi* (Eoc.-Georges Bank), 15(3) : 51, 9, 11.

Lagena acuticosta (Eoc.-Georges Bank), 15(3) : 59, 10, 32-34; (Eoc.-Ala.), 15(4) : 80, 14, 6; cf. *costata* (Eoc.-Georges Bank), 15(3) : 60, 10, 27; *costata* (Eoc.-Ala.), 15(4) : 80, 14, 7; *sulcata*, var. *apiculata* (Eoc.-Georges Bank), 15(3) : 60, 10, 28; sp.(?) (Eoc.-Georges Bank), 15(3) : 60, 10, 29-31.

Lamareckina torrei (Eoc.-Cuba), 13(1) : 21, 2, 24-26; *wilcoxensis* (Eoc.-Ala.), 15(4) : 84, 14, 30, 31.

Lenticulina convergens (Eoc.-Georges Bank), 15(3) : 54, 9, 24. *Lingulina cubensis* (Eoc.-Cuba), 13(1) : 11, 1, 38; (Eoc.-Georges Bank), 15(3) : 58, 10, 17; *taylorana* (Cret.-Texas), 14(2) : 43, 7, 9.

Listerella caribaea (Eoc.-Cuba), 13(1) : 6, 1, 29, 30; *cubana* (Eoc.-Cuba), 13(1) : 5, 1, 22, 23; *gracillima* (Eoc.-Cuba), 13(1) : 6, 1, 27, 28; *nodulosa*, var. *glabrata* (Tert.-Georges Bank), 11(4) : 77, 12, 1; *petrosa* (Eoc.-Cuba), 13(1) : 5, 1, 24-26.

Lituola irregulariter (Cret.-Texas), 15(4) : 89, 16, 9, 10.

[Louisiana]. New Species of Foraminifera from the Oligocene and Miocene, 15(1) : 1.

Loxostoma georgianum (Tert.-Georges Bank), 11(4) : 81, 12, 14, 15; *minutissimum* (Cret.-Texas), 14(2) : 45, 7, 19.

Marginulina, Some Notes on Cretaceous Species of, 13(4) : 91.

Marginulina texasensis Cushman, A New Name, 14(4) : 95.

Marginulina asperuliformis (Eoc.-Calif.), 15(2) : 24, 6, 1-3; *austinana* (Cret.-Texas), 13(4) : 92, 13, 1-4; var. *ascescens* (Cret.-Texas), 13(4) : 93, 13, 9; var. *directa* (Cret.-Texas), 13(4) : 93, 13, 5-8; cf. *bronni* (Cret.-Calif.), 11(3) : 67, 11, 4; *bullata* (Cret.-Texas), 13(4) : 96, 14, 9-15; *cocoaensis* (Eoc.-Georges Bank), 15(3) : 55, 9, 25; *cretacea* (Cret.-Texas), 13(4) : 94, 13, 12-15; *curvatura* (Cret.-Texas, Ark.), 14(2) : 34, 5, 13, 14; *dorsata* (Cret.-Texas), 13(4) : 94, 13, 16-18; *eximia* (Eoc.-Georges Bank), 15(3) : 56, 9, 27, 28; *georgiana* (Tert.-

- Georges Bank), 11(4): 77, 12, 2, 3; *gladius* (Eoc.-Georges Bank), 15(3): 54, 9, 32; *grata* (Cret.-Calif.), 11(3): 66, 10, 2; **havanensis** (Eoc.-Cuba), 13(1): 9, 1, 31, 32; **inconstantia** (Cret.-Texas), 14(2): 33, 5, 4-9; **jarvisi** (Cret.-Trinidad, Mexico), 14(2): 35, 5, 17, 18; **junceae** (Cret.-Texas), 14(2): 33, 5, 10; **laeviuscula** (Eoc.-Cuba), 13(1): 10, 1, 33, 34; **munda** (Cret.-Texas), 14(2): 34, 5, 11, 12; **navarroana** (Cret.-Ala., Texas), 13(4): 98, 14, 17, 18; **palmerae** (Eoc.-Cuba), 13(1): 10, 1, 36, 37; **plummerae** (Cret.-Texas), 13(4): 97, 13, 21-23; **pseudomarchi** (Cret.-Texas), 13(4): 94, 13, 19, 20; *silicula* (Cret.-Texas, Europe), 13(4): 97, 14, 19-22; **siliqua** (Cret.-Texas, Ark.), 14(2): 34, 5, 15, 16; **stephensoni** (Cret.-Texas), 13(4): 93, 13, 10, 11; **striato-carinata** (Cret.-Calif.), 11(3): 67, 10, 4, 7; **subbullata** (Eoc.-Georges Bank), 15(3): 55, 9, 30, 31; **taylorana** (Cret.-Texas), 13(4): 96, 14, 7; **tenuis** (Eoc.-Georges Bank), 15(3): 55, 9, 33, 34; **texana** (Cret.-Texas, Tenn.), 13(4): 95, 14, 1-4; **texasensis** (Cret.-Texas), 14(4): 95; *triangularis*, var. *panamensis* (Eoc.-Georges Bank), 15(3): 56, 9, 26; **trinitatensis** (Cret.-Trinidad), 13(4): 99, 14, 16; cf. *tripleura* (Cret.-Texas), 13(4): 95, 14, 5, 6; sp. (?) (Tert.-Calif.), 11(3): 58, 8, 12-14; (Eoc.-Georges Bank), 15(3): 56, 9, 29.
- Massilina decorata* (Eoc.-Georges Bank), 15(3): 52, 9, 13; **glabri-costata** (L. Olig.-Miss.), 11(2): 25, 4, 2; **texasensis** (Cret.-Texas), 13(4): 100, 15, 2, 3.
- [Miocene]. A New Genus of Foraminifera from the Miocene of Venezuela, 11(1): 13. Three New Foraminifera from the Miocene, Bowden Marl, of Jamaica, 12(1): 3. Notes on Some European Eocene [Miocene] Species of Bulimina, 13(2): 46. A New Species of Pulvinulinella from the California Miocene, 14(1): 1. New Species of Foraminifera from the Oligocene and Miocene, 15(1): 1. Notes on the Early Described Miocene Species of Uvigerina, 15(2): 33.
- [Mississippi]. Bitubulogenerina howei, A New Species from the Lower Oligocene, 11(1): 20. New Species of Foraminifera from the Lower Oligocene of Mississippi, 11(2): 25.
- Neobulimina, Notes on Some Cretaceous Species of Buliminella and, 12(1): 5.
- Neobulimina canadensis* (Cret.-Amer.), 12(1): 9, 2, 9, 10; **irregularis** (Cret.-Texas), 12(1): 9, 2, 8; **spinosa** (Cret.-Amer.), 12(1): 9, 2, 11.
- Neocarpenteria**, 12(2): 34; **cubana** (Eoc.-Cuba), 12(2): 34, 6, 3-5.

Nodobaculariella, Notes on Some of the Species Referred to Vertebralina and Articulina, and A New Genus, 13(2) : 41.

Nodobaculariella, 13(2) : 41; *atlantica* (Rec.-Atl.), 13(2) : 42, 5, 7, 8; *japonica* (Pleist.-Japan), 13(2) : 42, 5, 9-11.

Nodogenerina georgiana (Tert.-Georges Bank), 11(4) : 80, 12, 16; *havanensis* (Eoc.-Cuba), 13(1) : 14, 1, 47, 48; *laevis* (Eoc.-Cuba), 13(1) : 15, 2, 12; *sculpturata* (Eoc.-Georges Bank), 15(3) : 63, 10, 55.

Nodosarella acus (Eoc.-Cuba), 13(1) : 19, 2, 10, 11; *advena* (Eoc.-Calif.), 15(2) : 30, 6, 19, 20; *constricta* (Eoc.-Cuba), 13(1) : 18, 2, 4-7; *ignota* (Eoc.-Calif.), 15(2) : 30, 6, 21; *primitiva* (Cret.-Texas), 14(2) : 46, 8, 2; *subnodosa* (Eoc.-Georges Bank), 15(3) : 68, 11, 21, 22; *texana* (Cret.-Texas), 14(2) : 46, 8, 1; sp.(?) (Eoc.-Georges Bank), 15(3) : 68, 11, 23, 24.

Nodosaria cf. *alternata* (Cret.-Calif.), 11(3) : 71, 10, 8; cf. *anomala* (Tert.-Calif.), 11(3) : 58, 8, 16; *cookei* (Eoc.-Georges Bank), 15(3) : 58, 10, 12; *corsicanana* (Cret.-Texas), 14(2) : 42, 7, 1-4; *gracilitatis* (Cret.-Texas), 14(2) : 41, 6, 23-26; cf. *koina* (Tert.-Calif.), 11(3) : 58, 8, 15; *latejugata*, var. (Eoc.-Ala.), 15(4) : 80, 14, 4; *monile* (Cret.-Calif.), 11(3) : 71, 10, 5; *navarroana* (Cret.-Texas), 13(4) : 103, 15, 11; *spinifera* (Cret.-Calif.), 11(3) : 71, 10, 9, 10; *velascoensis* (Cret.-Calif.), 11(3) : 72, 11, 3; sp.(?) (Cret.-Calif.), 11(3) : 71, 11, 1, 2; sp.(?) B (Eoc.-Georges Bank), 15(3) : 58, 10, 13; sp.(?) A (Eoc.-Georges Bank), 15(3) : 58, 10, 14.

Nonion, Some New Species of, 12(3) : 63.

Nonion angustum (Plioc.-Cyprus), 12(3) : 68, 12, 11; *chapapotentse* (Eoc.-Georges Bank), 15(3) : 61, 10, 41-43; *decoratum* (Olig.-Ala.), 15(3) : 46, 9, 4; *dingdeni* (Mioc.-Germany), 12(3) : 65, 12, 5; *dollfusi* (Mioc.-France), 12(3) : 65, 12, 4; *halkyardi* (Eoc.-France), 12(3) : 63, 12, 1; *havanense* (Eoc.-Cuba), 13(1) : 19, 2, 13, 14; *ibericum* (Pleist.-Spain), 12(3) : 68, 12, 12, 13; *micrum* (Eoc.-Georges Bank), 15(3) : 61, 10, 44-46; *nicobarensis* (Plioc.-Kar Nicobar), 12(3) : 67, 12, 9; *novozealandicum* (Mioc.-New Zeal.), 12(3) : 66, 12, 6; *ornatissima* (Eoc.-Germany), 12(3) : 69, 12, 14; *planatum* (Eoc.-Ala.), 15(4) : 81, 14, 12, 13; *roemeri* (Olig.-Germany), 12(3) : 65, 12, 3; *schwageri* (Plioc.-Kar Nicobar), 12(3) : 67, 12, 8; *stachei* (Mioc.-New Zeal.), 12(3) : 66, 12, 7; *turgescens* (Olig.-Germany), 12(3) : 64, 12, 2; *victoriense* (Plioc.-Australia), 12(3) : 67, 12, 10; *wilcoxense* (Eoc.-Ala.), 15(4) : 81, 14, 14.

Nonionella ansata (Cret.-Ala.), 14(2) : 44, 7, 17; *crassipunctata*

(L. Olig.-Miss.), 11(2): 31, 4, 11; **frankei** (Olig.-Germany), 12(4): 88, 15, 10; **gardnerae** (Eoc.-Texas), 12(4): 88, 15, 9; **novo-zealandica** (Mioc.-New Zeal.), 12(4): 88, 13, 16; **oligocenica** (Olig.-Ala.), 15(3): 47, 9, 5; **parri** (Rec.-New Zeal.), 12(4): 89, 13, 17; **pauciloba** (L. Olig.-Miss.), 11(2): 32, 4, 12; **wilcoxensis** (Eoc.-Ala.), 15(4): 82, 14, 15.

[Oligocene]. *Bitubulogenerina howei*, A New Species from the Lower Oligocene, 11(1): 20. New Species of Foraminifera from the Lower Oligocene of Mississippi, 11(2): 25. Notes on Some Oligocene Species of Bulimina and Buliminella, 13(1): 36. Three New Rotaliform Foraminifera from the Lower Oligocene and Upper Eocene of Alabama, 14(3): 62. Notes on the Oligocene Species of Uvigerina and Angulogerina, 14(4): 74. New Species of Foraminifera from the Oligocene and Miocene, 15(1): 1. New Species of Foraminifera from the Lower Oligocene of Alabama, 15(3): 45.

[Pacific]. New Genera and Species of Foraminifera of the Late Tertiary of the Pacific, 12(2): 45.

Paleoecology as Shown by the Foraminifera, 15(2): 40.

Palmula pilulata (Cret.-Texas), 14(2): 37, 6, 2; **primitiva** (Cret.-Texas), 15(4): 91, 16, 4, 5; **simplex** (Cret.-Texas), 14(2): 36, 6, 1.

Planispirina mornhinvegi (L. Olig.-Miss.), 11(2): 27, 4, 6.

Planoglobulina, 14(1): 22; *acervulinoides* (Cret.-Europe, Amer.), 14(1): 23, 4, 5-8; *taylorana* (Cret.-Texas), 14(1): 23, 4, 9, 10.

Planularia sp. (Eoc.-Ala.), 15(4): 80, 13, 13.

Planulina austinana (Cret.-Texas), 14(3): 68, 12, 2; **harangensis** (Mioc.-La.), 15(1): 14, 2, 9; **nacatochensis** (Cret.-Ark.), 14(2): 50, 8, 9; **simplex** (Eoc.-Cuba), 13(1): 28, 2, 45-47; **spissocostata** (Cret.-Texas), 14(3): 69, 12, 4; **suturata** (Eoc.-Cuba), 13(1): 29, 2, 54-56; **texana** (Cret.-Texas), 14(3): 69, 12, 3.

Plectina cubensis (Eoc.-Cuba), 12(3): 57, 10, 7-9; **elongata** (Eoc.-Cuba), 12(3): 58, 10, 22-24; **garzaensis** (Eoc.-Calif.), 11(4): 92, 14, 3, 4; **torrei** (Eoc.-Cuba), 12(3): 57, 10, 12-14.

Plectofrondicularia angusti-costata (Tert.-Georges Bank), 11(4): 80, 12, 12; **basi-spinata** (Tert.-Georges Bank), 11(4): 79, 12, 11; **garzaensis** (Eoc.-Calif.), 15(2): 26, 6, 9; **georgiana** (Tert.-Georges Bank), 11(4): 79, 12, 13; cf. *miocenica* (Tert.-Calif.), 11(3): 59; **palmerae** (Eoc.-Cuba), 12(3): 61, 11, 20-24; **vaughani** (Tert.-Calif.), 11(3): 59, 9, 1.

[Pleistocene]. Notes on Some Pliocene and Pleistocene Species of Bulimina and Buliminella, 14(3) : 53.

Pleurostomella alazanensis, var. **cubensis** (Eoc.-Cuba), 13(1) : 17, 1, 64, 65; *bellardi* (Eoc.-Georges Bank), 15(3) : 67, 11, 16, 17; *cubensis* (Eoc.-Georges Bank), 15(3) : 67, 11, 25, 26; *jacksonensis* (Eoc.-Georges Bank), 15(3) : 67, 11, 20; *naranjoensis* (Eoc.-Cuba), 13(1) : 16, 1, 59, 60; *nuttalli* (Eoc.-Calif.), 15(2) : 29, 6, 17, 18; *obesa* (Eoc.-Cuba), 13(1) : 16, 1, 61; *rimosa* (Eoc.-Cuba), 13(1) : 17, 1, 62, 63; *tenuis* (Eoc.-Georges Bank), 15(3) : 67, 11, 18, 19.

[Pliocene]. Notes on Some Pliocene and Pleistocene Species of Bulimina and Buliminella, 14(3) : 53. *Rupertia* (?) *adamsi*, A New Species from the Pliocene of California, 15(2) : 21.

Polymorphinella, 12(2) : 46; **compressa** (Pleist.-Japan), 12(2) : 47, 8, 6; **vaginulinaeformis** (Pleist.-Japan), 12(2) : 47, 8, 5.

Polymorphinoides, 12(2) : 48; **spiralis** (Pleist.-Japan), 12(2) : 48, 8, 4.

Polystomellina australis (Rec.-Australia), 12(4) : 87, 15, 7; **miocenica** (Mioc.-Australia), 12(4) : 87, 15, 8.

Pseudoclavulina cocoaensis (Eoc.-Georges Bank), 15(3) : 50, 9, 17.

Pseudoglandulina gallowayi (Tert.-Calif.), 11(3) : 59, 8, 17; *laevigata* (Eoc.-Georges Bank), 15(3) : 58, 10, 15, 16; var. *ovata* (Eoc.-Ala.), 15(4) : 80, 14, 5.

Pseudopolymorphina wilcoxensis (Eoc.-Ala.), 15(4) : 81, 14, 11.

Pseudorbitolina cubensis (Eoc.-Cuba), 12(3) : 59, 10, 27-30.

Pseudotextularia, 14(1) : 21; *varians* (Cret.-Europe, Amer.), 14(1) : 21, 4, 1-4.

Pullenia americana (Cret.-Amer.), 12(4) : 76, 13, 4, 5; *bulloides* (Cret.-Europe), 12(4) : 74; *compressiuscula* (Cret.-Germany), 12(4) : 74; *coryelli* (Cret.-Mexico, Trinidad), 12(4) : 74; **cretacea** (Cret.-Amer.), 12(4) : 75, 13, 8; **eocenica** (Eoc.-Calif.), 15(2) : 31, 7, 1; *jarvisi* (Cret.-Mexico, Trinidad), 12(4) : 77, 13, 6; *minuta* (Cret.-Amer.), 12(4) : 77, 13, 7; *octoloba* (Tert.-Calif.), 11(3) : 63, 9, 9; *puentepiedraensis* (Cret.-Mexico), 12(4) : 75; *quaternaria* (Cret.-Germany), 12(4) : 74; *quinqueloba* (Cret.-Europe), 12(4) : 74; cf. *quinqueloba* (Eoc.-Georges Bank), 15(3) : 73, 12, 17; *sphaeroides* (Cret.-Germany), 12(4) : 74; cf. *sphaeroides* (Eoc.-Georges Bank), 15(3) : 73.

Pulvinulinella from the California Miocene, A New Species of, 14(1) : 1.

Pulvinulinella atlantisae (Eoc.-Georges Bank), 15(3) : 72, 12, 16;
cancellata (Eoc.-Cuba), 12(2) : 35, 6, 7; *culter*, var. *mexicana*
(Eoc.-Georges Bank), 15(3) : 72, 12, 15; **glabrata** (Cret.-
Texas), 14(3) : 66, 11, 4; **gyroidinaformis** (Mioc.-Calif.), 14(1) :
2, 1, 1, 2; **navarroana** (Cret.-Texas), 14(3) : 66, 11, 5; *obtusa*
(Eoc.-Ala.), 15(4) : 87, 15, 12, 13; **tenuicarinata** (Eoc.-Calif.),
11(4) : 95, 14, 11; **texana** (Cret.-Texas), 14(2) : 49, 8, 8.

Pyrgo oligocena (L. Olig.-Miss.), 11(2) : 26, 4, 5.

Pyrgoella, A New Genus of the Miliolidae, 12(4) : 90.

Pyrgoella, 12(4) : 90; *sphaera* (Rec.-Pac.), 12(4) : 90.

Pyrulina cf. *cylindroides* (Eoc.-Georges Bank), 15(3) : 61, 10, 37.

Quinqueloculina eocenica (Eoc.-Georges Bank), 15(3) : 51, 9, 12;
moremani (Cret.-Texas), 13(4) : 100, 15, 1; sp. (Eoc.-Ala.),
15(4) : 78, 13, 4.

Ramulina arkadelphiana (Cret.-Ark.), 14(2) : 43, 7, 12-14; **globo-**
tubulosa (Cret.-Tenn.), 14(2) : 44, 7, 16; **navarroana** (Cret.-
Texas), 14(2) : 43, 7, 10, 11; **ornata** (Cret.-Trinidad, Mexico),
14(2) : 44, 7, 15; sp.(?) (Eoc.-Georges Bank), 15(3) : 61, 10,
39, 40.

[Recent]. A New Genus of Foraminifera, *Discorbinella*, from
Monterey Bay, California, 11(4) : 89. The Recent Species of
Bulimina Named by d'Orbigny in 1826, 14(4) : 90.

Rectoeponides, 12(2) : 31; **cubensis** (Eoc.-Cuba), 12(2) : 31, 5, 18.

Rectogümbelina, 14(1) : 19; *cretacea* (Cret.-Ark.), 14(1) : 20, 3,
15, 16; **hispidula** (Cret.-Texas), 14(1) : 21, 3, 20-22; **minuta**
(Cret.-Texas), 14(2) : 45, 7, 18; *texana* (Cret.-Texas), 14(1) :
20, 3, 17-19.

Reophax cf. *pilulifer* (Tert.-Calif.), 11(3) : 55, 8, 2.

Robertina, Some Species of, 12(4) : 92. Two New Species of
Robertina, 14(4) : 73.

Robertina angusta (Olig.-Miss.), 12(4) : 96, 16, 11; *arctica* (Rec.-
Arctic), 12(4) : 93, 16, 1; *austriaca* (Mioc.-Austria), 12(4) : 94,
16, 2, 3; **bradyi** (Rec.-Atl.), 12(4) : 99, 16, 9; **californica** (Plioc.-
Calif.), 12(4) : 97, 16, 14; *charlottensis* (Rec.-E. Pac.), 12(4) :
97, 16, 12; *convoluta* (Rec.-E. Atl.), 12(4) : 94, 16, 4; *declivis*
(Olig.-Germany), 12(4) : 94, 16, 5; **germanica** (L. Olig.-Ger-
many), 14(4) : 73, 13, 2; *imperatrix* (Mioc.-Hungary), 12(4) :
95, 16, 6, 7; *ovigera* (Eoc.-France), 12(4) : 93, 16, 15; **plum-**
merae (Eoc.-Texas), 14(4) : 73, 13, 1; *subcylindrica* (Rec.-
Atl.), 12(4) : 95, 16, 10; **translucens** (Rec.-Atl.), 12(4) : 99, 16,

- 8; *wilcoxensis* (Eoc.-Ala.), 12(4): 96, 16, 13; (Eoc.-Ala.), 15(4): 82, 14, 16.
- Robulus atlantisae* (Eoc.-Georges Bank), 15(3): 54, 9, 22, 23; *barbati* (Tert.-Calif.), 11(3): 57, 8, 9; *limbosus* (Eoc.-Georges Bank), 15(3): 52, 9, 20; var. *hockleyensis* (Eoc.-Georges Bank), 15(3): 53, 9, 21; *mayeri* (Mioc.-La.), 15(1): 4, 1, 6; cf. *mayi* (Tert.-Calif.), 11(3): 57, 8, 10; *navarroensis*, var. *extruatus* (Cret.-Texas), 14(2): 31, 5, 1; *pseudo-secans* (Cret.-Tenn.), 14(2): 32, 5, 3; *spisso-costatus* (Cret.-Texas), 14(2): 32, 5, 2; *stephensoni* (Cret.-Tenn.), 15(4): 90, 16, 2, 3; *submarinus* (Eoc.-Georges Bank), 15(3): 53, 9, 18, 19; *warmani* (Tert.-Calif.), 11(3): 56, 8, 8; sp. (?) (Tert.-Calif.), 11(3): 56, 8, 7; (Eoc.-Ala.), 15(4): 79, 13, 9, 10.
- Rupertia* (?) *adamsi*, A New Species from the Pliocene of California, 15(2): 21.
- Rupertia* (?) *adamsi* (Tert.-Calif.), 15(2): 22, 5, 1-11.
- Saracenaria ornatula* (Eoc.-Cuba), 13(1): 11, 1, 35; *schlencki* (Tert.-Calif.), 11(3): 57, 8, 11.
- Sigmomorphina cubensis* (Eoc.-Cuba), 13(1): 12, 1, 41; *transversa*, var. *cubensis* (Eoc.-Cuba), 13(1): 16, 2, 8, 9.
- Siphogenerinoides* from California, A New, 12(4): 91.
- Siphogenerinoides brevispinosa* (Cret.-Ark.), 15(4): 92, 16, 7; *clarki* (Cret.-Calif.), 12(4): 91, 13, 9-12; 13(2): 53.
- Siphonina advena*, var. *cubensis* (Eoc.-Cuba), 13(1): 24, 2, 19, 20; *davisi* (Olig.-La.), 15(1): 10, 2, 3; *nuda* (Eoc.-Cuba), 12(3): 62, 11, 3-6; *pustulata* (Eoc.-Cuba), 12(3): 62, 11, 1, 2; *wilcoxensis* (Eoc.-Ala.), 15(4): 86, 15, 7-9.
- Sphaeroidina bulloides* (Cret.-Europe, Australia), 12(4): 78.
- Spirillina selseyensis* (Eoc.-Ala.), 15(4): 84, 14, 28, 29; *vicksburgensis* (L. Olig.-Miss.), 11(2): 34, 5, 6.
- Spirolina arrecta* (L. Olig.-Miss.), 11(2): 27, 4, 7.
- Spiroloculina wilcoxensis* (Eoc.-Ala.), 15(4): 78, 13, 5, 6; sp. (Eoc.-Ala.), 15(4): 79, 13, 7.
- Spiroplectamina adamsi* (Tert.-Calif.), 11(2): 39, 6, 1, 2; *alexanderi* (L. Cret.-Texas), 11(1): 1, 1, 1; *barrowi* (Mioc.-La.), 15(1): 1, 1, 1; *chicoana* (Cret.-Calif.), 11(1): 7, 1, 8, 9; *gaultana* (L. Cret.-England), 11(1): 8, 2, 1, 2; *goodlandana* (L. Cret.-Texas), 11(1): 2, 1, 2, 3; *jarvisi* (Cret.-Trinidad), 15(4): 90, 16, 1; *longa* (L. Cret.-Texas), 11(1): 3, 1, 4, 5; *mexiaensis* (Eoc.-Texas), 11(2): 43, 6, 5, 6; *nuda* (L. Cret.-Texas), 11(1): 4, 1, 6, 7; *nuttalli* (Eoc.-Venezuela), 11(2): 43, 6, 3, 4; *roemeri* (Cret.-Germany), 11(1): 9, 2, 5; *senonana*

(Cret.-Germany), 11(1): 9, 2, 3, 4; *sigmoidina* (Cret.-Calif.), 11(1): 7, 1, 10, 11; *wilcoxensis* (Eoc.-Ala.), 15(4): 78, 13, 1, 2. *Spiroplectoides attenuata* (Eoc.-Georges Bank), 15(3): 62, 10, 47-49; *cubensis* (Eoc.-Cuba), 13(1): 13, 1, 44, 45; *directa* (Eoc.-Calif.), 15(2): 26, 6, 7, 8. *Stichocibicides*, 12(2): 33; *cubensis* (Eoc.-Cuba), 12(2): 33, 5, 19-21.

[Taxonomic]. New Cretaceous Textulariidae, 11(1): 1. A New Genus of Foraminifera from the Miocene of Venezuela, 11(1): 13. New Late Tertiary Bolivinas from California, 11(1): 16. *Bitubulogenerina howei*, A New Species from the Lower Oligocene, 11(1): 20. New Species of Foraminifera from the Lower Oligocene of Mississippi, 11(2): 25. New Tertiary Textulariidae, 11(2): 39. A Foraminiferal Faunule from the Type San Lorenzo Formation, Santa Cruz County, California, 11(3): 53. Cretaceous Foraminifera from the Moreno Shale of California, 11(3): 65. Some New Foraminifera from the Late Tertiary of Georges Bank, 11(4): 77. Notes on Some American Cretaceous Flabellinas, 11(4): 83. A New Genus of Foraminifera, *Discorbinella*, from Monterey Bay, California, 11(4): 89. New Species of Foraminifera from the Kreyenhagen Shale of Fresno County, California, 11(4): 90. Some American Cretaceous Buliminas, 11(4): 96. The Foraminiferal Genus *Amphimorphina* in the Eocene of Cuba, 12(1): 1. Three New Foraminifera from the Miocene, Bowden Marl, of Jamaica, 12(1): 3. Notes on Some Cretaceous Species of *Buliminella* and *Neobulimina*, 12(1): 5. Notes on Some American Cretaceous *Fronicularias*, 12(1): 11. New Genera and Species of Foraminifera from the Eocene of Cuba, 12(2): 27. Some American Eocene Buliminas, 12(2): 39. New Genera and Species of Foraminifera of the Late Tertiary of the Pacific, 12(2): 45. Some American Cretaceous Species of *Ellipsonodosaria* and *Chrysalogonium*, 12(3): 51. Additional New Species of Foraminifera and A New Genus from the Eocene of Cuba, 12(3): 55. Some New Species of *Nonion*, 12(3): 63. Cretaceous Foraminifera of the Family *Chilostomellidae*, 12(4): 71. Some New Species of *Elphidium* and Related Genera, 12(4): 78. *Pyrgoella*, A New Genus of the *Miliolidae*, 12(4): 90. A New *Siphogenerinoides* from California, 12(4): 91. Some Species of *Robertina*, 12(4): 92. Further New Species of Foraminifera from the Eocene of Cuba, 13(1): 1. *Astrononion*, A New Genus

of the Foraminifera, and Its Species, 13(1): 29. Notes on Some Oligocene Species of Bulimina and Buliminella, 13(1): 36. Notes on Some of the Species Referred to Vertebralina and Articulina, and A New Genus Nodobaculariella, 13(2): 41. Notes on Some European Eocene [Miocene] Species of Bulimina, 13(2): 46. Notes on the Early Described Eocene Species of Uvigerina and Some New Species, 13(2): 54. Notes on Some of the Early Described Eocene Species of Bulimina and Buliminella, 13(3): 65. The Described American Eocene Species of Uvigerina, 13(3): 74. Some Notes on Cretaceous Species of Marginulina, 13(4): 91. A Few New Species of American Cretaceous Foraminifera, 13(4): 100. Additional New Species of Eocene Foraminifera from Cuba, 13(4): 106. A New Species of Pulvinulinella from the California Miocene, 14(1): 1. Cretaceous Species of Gümbelina and Related Genera, 14(1): 2. Some New Names in the Foraminifera, 14(1): 28. Additional New Species of American Cretaceous Foraminifera, 14(2): 31. Notes on Some Pliocene and Pleistocene Species of Bulimina and Buliminella, 14(3): 53. Three New Rotaliform Foraminifera from the Lower Oligocene and Upper Eocene of Alabama, 14(3): 62. Some New Species of Rotaliform Foraminifera from the American Cretaceous, 14(3): 66. Two New Species of Robertina, 14(4): 73. Notes on the Oligocene Species of Uvigerina and Angulogerina, 14(4): 74. The Recent Species of Bulimina Named by d'Orbigny in 1826, 14(4): 90. Marginulina texasensis Cushman, A New Name, 14(4): 95. New Species of Foraminifera from the Oligocene and Miocene, 15(1): 1. Cribrolinoides, A New Genus of the Foraminifera, Its Development and Relationships, 15(1): 15. Rupertia (?) adamsi, A New Species from the Pliocene of California, 15(2): 21. Some New and Interesting Foraminifera from the Kreyenhagen Shale of California, 15(2): 23. Notes on the Early Described Miocene Species of Uvigerina, 15(2): 33. New Species of Foraminifera from the Lower Oligocene of Alabama, 15(3): 45. Eocene Foraminifera from Submarine Cores off the Eastern Coast of North America, 15(3): 49. Eocene Foraminifera of Wilcox Age from Woods Bluff, Alabama, 15(4): 77. New American Cretaceous Foraminifera, 15(4): 89. Bulimina macilenta Cushman and Parker, A New Name, 15(4): 93. [Tertiary]. New Late Tertiary Bolivinas from California, 11(1): 16. A Foraminiferal Faunule from the Type San Lorenzo Formation, Santa Cruz County, California, 11(3): 53. Some

New Foraminifera from the Late Tertiary of Georges Bank, 11(4): 77. New Tertiary Textulariidae, 11(2): 39. New Genera and Species of Foraminifera of the Late Tertiary of the Pacific, 12(2): 45.

Textularia americana, 14(1): 5, 1, 19, 20; *ampliata*, 14(1): 5, 1, 14; *euryconus*, 14(1): 6, 1, 23; *globifera*, 14(1): 6, 1, 26, 27; *globulosa*, var. *amplior*, 14(1): 5, 1, 9; var. *dilatata*, 14(1): 5, 1, 10; var. *obtusa*, 14(1): 5, 1, 8; *gomphoconus*, 14(1): 6, 1, 24; *inflata*, 14(1): 5, 1, 12; *leptotheca*, 14(1): 5, 1, 15; *missouriensis*, 14(1): 6, 1, 21; *pachyaulax*, 14(1): 5, 1, 16-18; *poroconus*, 14(1): 6, 1, 22; *subtilis*, 14(1): 5, 1, 11; *sulcata*, 14(1): 5, 1, 13.

Textularia aciculata, 14(1): 3; *anglica* (L. Cret.-England), 11(1): 10, 2, 6, 7; *aspera* 14(1): 4, 1, 3, 4; *badenensis* (Mioc.-Austria), 11(2): 44, 7, 1; *bartonana* (Eoc.-Italy), 11(2): 44, 7, 6; *chapmani* (L. Cret.-England), 11(1): 13, 2, 8, 9; *dilatata*, 14(1): 4, 1, 6; *dollfussi* (Mioc.-France), 11(2): 45, 7, 8, 9; *globulosa*, 14(1): 4; cf. *gramen* (Tert.-Calif.), 11(3): 56, 8, 5; *halkyardi* (Eoc.-France), 11(2): 45, 7, 5; *laevis*, 14(1): 4, 1, 5; *lajollensis* (Eoc.-Calif.), 11(2): 46, 6, 13; *lontensis* (Eoc.-Italy), 11(2): 49, 7, 2, 3; *midwayana* (Eoc.-Texas), 11(2): 49, 6, 7-9; *ovulata* (Mioc.-France), 11(2): 50, 6, 11, 12; *perforata*, 14(1): 4, 1, 7; *plummerae* (Eoc.-Texas), 11(2): 50, 6, 10; *spinosa*, 14(1): 4; *striata*, 14(1): 4; *suttonensis* (Plioc.-England), 11(2): 51, 7, 4; *tatumi* (Mioc.-La.), 15(1): 2, 1, 2; *teasi* (Olig.-La.), 15(1): 2, 1, 3; *thanetana* (Eoc.-England), 11(2): 51, 7, 7.

Trifarina advena (Eoc.-Georges Bank), 15(3): 66, 11, 15.

Triloculina mississippiensis (L. Olig.-Miss.), 11(2): 25, 4, 1.

Tritaxilina colei (Eoc.-Calif.), 11(4): 92, 14, 5, 6; *cubensis* (Eoc.-Cuba), 13(1): 7.

Trochammina taylorana (Cret.-Texas), 13(4): 101, 15, 4.

Unicosiphonia, 11(4): 81; *crenulata* (Tert.-Georges Bank), 11(4): 81, 12, 9, 10.

Uvigerina and Some New Species, Notes on the Early Described Eocene Species of, 13(2): 54. The Described American Eocene Species of *Uvigerina*, 13(3): 74. Notes on the Oligocene Species of *Uvigerina* and *Angulogerina*, 14(4): 74. Notes on the Early Described Miocene Species of *Uvigerina*, 15(2): 33.

Uvigerina abbreviata (Eoc.-France), 13(2): 56, 8, 6, 7; *aculeata* (Mioc.-Austria), 15(2): 35, 8, 1-5; (Mioc.-Germany), 15(2): 37; *alabamensis* (Eoc.-Ala.), 15(4): 83, 14, 26, 27; *alata* (Eoc.-

Texas), 13(3) : 74, 11, 1, 2; *alazanensis* (Olig.-Mexico), 14(4) : 77, 13, 17; *altacostata* (Mioc.-La.), 15(1) : 9, 1, 14; *asperula* (Mioc.-Austria), 15(2) : 36; *aubेरiana* (Olig.-Venezuela), 14(4) : 89, 14, 13; (Olig.-Mexico), 14(4) : 89, 14, 18; *beccarii* (Olig.-Ecuador), 14(4) : 75, 13, 7; *biserialis* (Eoc.-France), 13(2) : 59, 8, 11, 12; *blanca-costata* (Eoc.-Mexico), 13(3) : 74, 11, 3, 4; *bonairensis* (Eoc.-Bonaire), 13(3) : 75, 11, 5; *boqueronensis* (Eoc.-Panama), 13(3) : 75, 11, 6; *brunnensis* (Mioc.-Austria), 15(2) : 37; *camagueyana* (Eoc.-Cuba), 13(3) : 76, 11, 7; *canariensis* (Eoc.-Peru), 13(3) : 87; var. *spinulosa* (Olig.-Cuba), 14(4) : 78, 14, 9; *capayana* (Olig.-Venezuela), 14(4) : 80, 14, 1; *carapitana* (Olig.-Cuba), 14(4) : 82, 14, 2; *churchi* (Eoc.-Calif.), 15(2) : 29, 6, 16; *cochlearis* (Mioc.-Austria), 15(2) : 36; *cocoaensis* (Eoc.-Ala., Miss.), 13(3) : 76, 11, 8, 9; (Olig.-Oregon), 14(4) : 89, 14, 15-17; *cookei* (Eoc.-Amer.), 13(3) : 76, 11, 10, 11; *cubana* (Olig.-Cuba), 14(4) : 79, 13, 18, 19; *curta* (Eoc.-Trinidad), 13(3) : 77, 11, 12, 13; *danvillensis* (Eoc.-La.), 13(3) : 77, 11, 14; *dumblei* (Eoc.-Texas), 13(3) : 78, 11, 17; *ecuadorensis* (Olig.-Ecuador), 14(4) : 83, 14, 7; *elongata* (Eoc.-Mexico), 13(3) : 78, 11, 15, 16; (Eoc.-Georges Bank), 15(3) : 65, 11, 11; *eocaena* (Eoc.-Germany), 13(2) : 54, 8, 1, 2; *farinosa* (Eoc.-Hungary), 13(2) : 58, 8, 8, 9; (Eoc.-La.), 13(3) : 87; *fragilis* (Eoc.-France), 13(2) : 57; *gallowayi* (Olig.-Ecuador), 14(4) : 75, 13, 8, 9; *gardnerae* (Eoc.-Amer.), 13(3) : 79, 11, 19, 20; var. *cubana* (Olig.-Cuba), 14(4) : 79, 13, 13; var. *texana* (Eoc.-Texas, La.), 13(3) : 79, 11, 18; *garzaensis* (Eoc.-Calif.), 15(2) : 28, 6, 15; *glabrans* (Eoc.-Ala., So. Car.), 13(3) : 80, 11, 21; *gladysae* (Eoc.-Mexico), 13(3) : 80, 12, 1; *globosa* (Mioc.-Phil. Ids.), 15(2) : 37; *gracilis* (Olig.-Germany), 14(4) : 74, 13, 3-6; *hantkeni* (Eoc.-Hungary), 13(2) : 60, 8, 15, 16; *havanensis* (Eoc.-Cuba), 12(3) : 59, 10, 19-21; (Eoc.-Cuba), 13(3) : 80, 12, 2, 3; *jacksonensis* (Eoc.-Amer.), 13(3) : 81, 12, 4, 5; *lappa* (Eoc.-Austria), 13(2) : 58, 8, 10; *laubeana* (Mioc.-Europe), 15(2) : 38; *liretensis* (Mioc.-La.), 15(1) : 7, 1, 13; *longa* (Eoc.-Cuba), 13(4) : 109, 16, 5, 6; var. *denticosta* (Eoc.-Cuba), 13(4) : 109, 16, 2, 3; *mantaensis* (Olig.-Ecuador), 14(4) : 84, 14, 8; *mariannensis* (Olig.-Fla.), 14(4) : 76, 13, 12; *mediterranea* (Eoc.-Bonaire), 13(3) : 87; *mexicana* (Olig.-Mexico), 14(4) : 77, 13, 14, 15; *neudorfensis* (Mioc.-Hungary), 15(2) : 39; *nuda* (Eoc.-France), 13(2) : 57; *nuttalli* (Olig.-Mexico), 14(4) : 82, 14, 3-5; *orbignyana* (Mioc.-Austria), 15(2) : 36; *parkeri* (Mioc.-Austria), 15(2) : 37, 8, 6, 7; *pigmea* (Eoc.-Amer.),

13(3) : 87; (Olig.-Mexico), 14(4) : 89, 14, 14; *pilulata* (Olig.-La.), 15(1) : 7, 1, 12; *rippensis* (Eoc.-Mexico), 13(3) : 81, 12, 6; (Eoc.-Georges Bank), 15(3) : 66, 11, 13, 14; *rugosa* (Eoc.-France), 13(2) : 58; *rustica* (Olig.-Ecuador), 14(4) : 83, 14, 6; *semiornata* (Olig.-France), 14(4) : 89, 14, 11, 12; (Mioc.-Austria), 15(2) : 34, 8, 8-14; *semivestita* (Eoc.-Cuba), 13(3) : 82, 12, 7, 8; *seriata* (Eoc.-Trinidad), 13(3) : 82, 12, 9, 10; *spini-costata* (Eoc.-Trinidad), 13(3) : 83, 12, 11, 12; (Eoc.-Georges Bank), 15(3) : 66, 11, 12; var. *alazanensis* (Olig.-Mexico), 14(4) : 78, 13, 16; *spinulosa* (Eoc.-Panama), 13(3) : 83, 12, 13; *tenuistriata* (Olig.-Mexico), 14(4) : 89, 14, 19; *topilensis* (Eoc.-Mexico), 13(3) : 83, 12, 14; *trilobata* (Mioc.-France), 15(2) : 34; *uhligi* (Mioc.-Europe), 15(2) : 38; *urnula* (Mioc.-Austria), 15(2) : 34, 8, 19-26; *venusta* (Mioc.-Hungary), 15(2) : 38, 8, 15-18; *vicksburgensis* (Olig.-Texas), 14(4) : 76, 13, 10, 11; *westermanni* (Eoc.-Bonaire), 13(3) : 84, 12, 15; *wilcoxensis* (Eoc.-Ala.), 15(4) : 82, 14, 17, 18; *yazooensis* (Eoc.-Amer.), 13(3) : 85, 12, 16, 17; *yeguaensis* (Eoc.-Texas), 13(3) : 86, 12, 18; sp.(?) (Olig.-Germany), 14(4) : 84, 14, 10.

Uvigerinella obesa var. *impolita* (Tert.-Calif.), 11(3) : 62, 9, 6.

Vaginulina atlantisae (Eoc.-Georges Bank), 15(3) : 59, 10, 23-26; *deprofunda* (Eoc.-Georges Bank), 15(3) : 58, 10, 18-22; *selmaensis* (Cret.-Tenn.), 13(4) : 101, 15, 8, 9; cf. *simondsi* (Cret.-Calif.), 11(3) : 69, 11, 7; *subgracilis* (Cret.-Texas), 13(4) : 103, 15, 13; *suturalis* (Cret.-Ark.), 13(4) : 102, 15, 5-7; *taylorana* (Cret.-Texas), 14(2) : 36, 5, 19.

Valvulammina affinis (Eoc.-Cuba), 13(1) : 9; (Eoc.-Cuba), 13(2) : 5, 3; *cubensis* (Eoc.-Cuba), 13(1) : 7; (Eoc.-Cuba), 13(2) : 5, 2.

Valvulina martii (Eoc.-Cuba), 13(1) : 7; (Eoc.-Cuba), 13(2) : 5, 1.

Valvulineria advena (Eoc.-Calif.), 15(2) : 31, 6, 22; *cubana* (Eoc.-Cuba), 12(2) : 34, 6, 1, 2; *georgiana* (Tert.-Georges Bank), 11(4) : 82, 12, 17; *paucilocula* (L. Olig.-Miss.), 11(2) : 37, 5, 7; *scrobiculata* (Eoc.-Ala.), 15(4) : 85, 14, 32, 33; *sculpturata* (L. Olig.-Miss.), 11(2) : 37, 5, 10; *wilcoxensis* (Eoc.-Ala.), 15(4) : 85, 15, 1, 2; sp.(?) (Eoc.-Georges Bank), 15(3) : 71.

[Venezuela]. A New Genus of Foraminifera from the Miocene of Venezuela, 11(1) : 13.

Ventilabrella, 14(1) : 24; *austinana* (Cret.-Amer.), 14(1) : 26, 4, 19; *carseyae* (Cret.-Amer.), 14(1) : 26, 4, 20-24; *decurrens* (Cret.-England), 14(1) : 27, 4, 18; *eggeri* (Cret.-Europe,

- Amer.), 14(1): 25, 4, 12-14; var. *glabrata* (Cret.-Texas), 14(1): 26, 4, 15-17; *ornatissima* (Cret.-Calif.), 14(1): 27, 4, 11.
- Verneuilina ornata* (Cret.-Europe), 14(1): 28; *pinarensis* (Eoc.-Cuba), 13(1): 1, 1, 3, 4; *villarensis* (Eoc.-Cuba), 13(1): 1, 1, 1, 2; sp.(?) (Tert.-Calif.), 11(3): 56, 8, 6.
- Vertebralina and Articulina, and A New Genus Nodobaculariella, Notes on Some of the Species Referred to, 13(2): 41.
- Vertebralina*, 13(2): 43; *advena* (Olig.-Miss.), 13(2): 44; *cassis* (Rec.-Atl.), 13(2): 44; *contracta* (Eoc.-France), 13(2): 44; *elongata* (Mioc.-Hungary), 13(2): 44; *insignis* (Rec.-Pac.), 13(2): 44; *jamaicensis* (Eoc.-Jamaica), 13(2): 44; *laevigata* (Eoc.-France), 13(2): 44; *mucronata* (Rec.-Atl.), 13(2): 44; *sarmatica* (Mioc.-Austria), 13(2): 44; *striata* (Rec.-Atl.), 13(2): 43, 5, 4-6.
- Virgulina alabamensis* (Olig.-Ala.), 15(3): 48, 9, 7; *cylindrica* (Eoc.-Cuba), 12(2): 30, 5, 15; *exilis* (Mioc.-La.), 15(1): 6, 1, 11; *miniacea* (Eoc.-Cuba), 12(2): 30, 5, 14; *wilcoxensis* (Eoc.-Ala.), 15(4): 82, 14, 19-21.
- Vulvulina advena* (Eoc.-Georges Bank), 15(3): 50, 9, 9, 10; *curta* (Eoc.-Calif.), 11(4): 91, 14, 1, 2.

